Advancing Lunar & Solar System Science & Exploration Through a Lunar Sample Return Campaign

Clive R. Neal¹ & Sam Lawrence²

Lunar Exploration Analysis Group


neal.1@nd.edu, samuel.j.lawrence@nasa.gov


1 = University of Notre Dame2 = NASA Johnson Space Center



We need more samples than Apollo, Luna, & Lunar Meteorites

Subsequent missions have shown that the sample return sites were not ideal for exploring the Moon.






Lunar Terranes


- Apollo sites close to terrane boundaries.
- Samples contain PKT signature.

Jolliff et al. (2000) JGR 105, 4197


 Apollo sample collection is not representative of the lunar compositional diversity.

Giguere et al. (2000) MaPS 35, 193






New Lunar Lithologies




How do these new Rock Types (not represented in the sample collection) revise models of lunar evolution?

Pure Anorthosite or PAN: Kaguya



Yamamoto et al. (2012) *GRL* **39**, L13201, doi:10.1029/2012GL052098

New Lunar Lithologies




Chandrayaan-1: M3. Moscoviense Basin

How do these new Rock Types (not represented in the sample collection) revise models of lunar evolution?

Olivine, Orthopyroxene, and Mg-Spinel-rich lithologies ("OOS")

Pieters et al. (2011) JGR, 116, E00G08


Olivine-rich mantle (?) deposits



Yamamoto et al. (2010) *Nature Geosci.* **3**, 533-536.

Yamamoto et al. (2012) *Icarus* **218**, 331-344.

Non-Mare Silicic Magmatism



Gruithuisen
Domes

LaHire
LaHire
La, 2000|
Apennine
Bench

Lassell
Massif

Helmet, Montes
Riphaeus, Darney

- Enhanced Th;
- Explosive volcanism;
- Christensen frequency (CF)
 value (DIVINER) indicates
 silica-rich lithologies.



Kusuma et al. (2012) Planet. Space Sci. 67, 46-56.

Jolliff B.L. et al. (2011) Nature Geosci. 4, 566-571.


Glotch T.D. et al. (2011) GRL 111, L21204, doi:10.1029/2011GL049548.


Glotch T.D. et al. (2010) Science 329, 1510-1513

Hagerty et al. (2006) JGR 111, E06002, doi:10.1029/2005JE002592.

Lawrence et al. (2005) GRL. 32, L07201, doi:10.1029/2004GL022022...


Hawke et al. (2003) JGR 108 (E7), 5069, doi: 10.1029/2002JE002013.





Recent Volcanic Activity

"Recent" volcanic eruptions ~ 1 Ga.




Hiesinger et al. (2003) JGR 108, (doi: 10.1029/2002JE001985)

Hiesinger et al. (2010) JGR 115, E03003, doi:10.1029/2009JE003380

Hiesinger et al. (2011) GSA Spec. Pap. 477, 1-52

Recent Volcanic Activity





Ina depression in Lacus Felicitatis

Schultz et al. (2006) Nature, 444, 184-186.

- What are the ages of the IMPs?
- What are the source regions for these potentially "young" basalts?
- Implications for the thermal history of the Moon?
- What are the mechanisms of eruption?


Unname	ed .	Unnamed
32	33	38
4 - A.M.	₹ - K ह	
Unnamed	Unnamed	Carrel-1

to be older (~3.5 Ga):


- Head & Wilson (2017) Icarus 183, 176-223.
- Qiao et al. (2017) Geology,

Images = 450 m across Braden S.E. et al. (2014) Nature Geoscience, v. 7, p. 787-791.

Cratering Chronology



- Constraining crater chronology.
- Important for Solar System Science.
- Need unambiguous impact melt
- Was there a "cataclysm" around 3.9 Ga?
- What are the ages/fluxes of the older and younger ends of the crater count curve?



Copernicus

Teielkovekii SPA

Mare Trigenii Apollo

Teielkovekii SPA

Mare Australe

Schrodinger

Lyet Bauthy

SOUTH POLE


Barlow (2010) GSA Bull. 122, 644-657

Pyroclastic Deposits

http://astrogeology.usgs.gov/geology/moon-pyroclastic-volcanism-project


Lunar Pyroclastic Deposits: Locations of all deposits overlaid on Clementine 750-nm global mosaic, Simple Collection replaction (context of C

- Do different pyroclastic deposits of different ages indicate a consistent volatile composition of the lunar mantle?
- Can these be used as a resource for human exploration?




000 Kilometers

LunPyroDb51999.shp Global30grid.shp




Farside Highlands

Crustal thickness dichotomy (CoM offset from CoF)



Wieczorek et al. (2013) *Science* **339**, 671-675.

How did the lunar crust form?



Ohtake et al. (2012) Nature Geoscience 5, 384-388.


LETTERS

DUBLISHED ONLINE-29 APPLI 2012 | DOI-101038/NGE01458

nature geoscience

Asymmetric crustal growth on the Moon indicated by primitive farside highland materials

Makiko Ohtake¹*, Hiroshi Takeda², Tsuneo Matsunaga³, Yasuhiro Yokota³, Junichi Haruyama¹, Tomokatsu Morota⁴, Satoru Yamamoto³, Yoshiko Ogawa⁵, Takahiro Hiroi⁶, Yuzuru Karouji⁷, Kazuto Saiki⁸ and Paul G. Lucey⁹





Ice Permatrost tround PSRs

Neutron Suppression Regions (NSRs) are found in both Permanently Shadowed Regions (PSRs) and illuminated areas, and they are not coincident

with PSRs.

Possible with nearside sample return?






Colaprete et al. (2010) Science 330, 463-468. Colaprete et al. (2012) Space Sci. Rev. 167, 3-22

Future Sample Return Missions Sample Return:

- New lithologies, including potential mantle samples.
- South Pole-Aitken Basin impact melt ("MoonRise").
- Other younger (e.g., Copernicus, Tycho) impact craters.
- Multi-ring basins (Nectaris, Imbrium, and Orientale).
- "Young" volcanic features (e.g., Ina Structure).
- Felsic domes (Gruithuisen Domes, Hansteen-Alpha, Compton-Belkovich).
- Large pyroclastic deposits.
- Cryogenic sample return PSRs.




New
Lithologies:
Spinel




New Lithologies: Mantle?

Olivine-rich




New Lithologies: Mantle?




New Lithologies: PAN

PAN




Craters NE of Hevesy Schwarzschild Birkoff Compton Belkovich. Nevy Moscoviense Lithologies: Mare Smythii 5:77 Mendeleev Korolev Tsiolkovsky Ingenii Oppenheimer Chebyshev Leibnitz




★ Felsic
Felsic
Igneous
Complexes






Pyroclastic Deposits



Pyroclastic Deposits







Young Igneous








Hydrogen Deposits





Cabeus Shoemaker/Faustini/Nobile

Hydrogen Deposits



Targeted Sample Return - farside





Technology Development Robotic SR:

- Landers and sample return vehicles.
- Cryogenic sampling, transport, and curation.
- Rover development to survive the lunar day/night/PSRs temperature swings, sample identification, collection, and storage (including cryogenic capabilities).
- Development of a Moon Assent Vehicle to return the samples.
- These developments would have feed forward implications for Mars sample return or SR from other destinations.

Conclusions

- Private companies are developing lunar surface exploration capabilities, including sample return allowing regular access to Moon.
- Since a regular cadence of missions to the Moon would be required for private commercial companies to build a business case, we have an opportunity to change the paradigm of planetary science and exploration and implement an affordable lunar robotic program.
- A dedicated Lunar Science & Exploration Program (LSEP) Office could be established that involves the lunar community and industrial partners in mission planning and flight opportunities.
- A focused lunar program would allow NASA to be a regular customer while developing new capabilities and implementing at least some of the objectives listed in the Decadal Survey, the LEAG Exploration Roadmap, the SEM Report, etc.